

Quick Reference Guide HEDIS® MY 2023

FOR MORE INFORMATION, VISIT WWW.NCQA.ORG









Marketplace



Medicare

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA). The HEDIS measures and specifications were developed by and are owned by NCQA. NCQA holds a copyright in these materials and may rescind or alter these materials at any time. Users of the HEDIS measures and specifications shall not have the right to alter, enhance, or otherwise modify the HEDIS measures and specifications, and shall not disassemble, recompile, or reverse engineer the HEDIS measures and specifications. Anyone desiring to use or reproduce the materials, subject to licensed user restrictions, without modification for an internal non-commercial purpose may do so without obtaining any approval from NCQA. Use of the Rules for Allowable Adjustments of HEDIS to make permitted adjustments of the materials does not constitute a modification. All other uses, including a commercial use, or any external reproduction, distribution, or publication must be approved by NCQA and are subject to a license at the discretion of NCQA.

HEDIS® MY 2023 Quick Reference Guide

Updated to reflect NCQA HEDIS® MY 2023 Technical Specifications

NH Healthy Families, Ambetter from NH Healthy Families and Wellcare strives to provide quality healthcare to our membership as measured through HEDIS® quality metrics. We created the HEDIS® MY 2023 Quick Reference Guide to help you increase your practice's HEDIS® rates and address care opportunities for your patients. Please always follow the state and/or CMS billing guidance and ensure the HEDIS® codes are covered prior to submission.



What is HEDIS®?

HEDIS® (Healthcare Effectiveness Data and Information Set) is a set of standardized performance measures developed by the National Committee for Quality Assurance (NCQA) to objectively measure, report, and compare quality across health plans. NCQA develops HEDIS® measures through a committee represented by purchasers, consumers, health plans, health care providers, and policy makers.



What are the scores used for?

As state and federal governments move toward a quality-driven healthcare industry, HEDIS® rates are becoming more important for both health plans and individual providers. State purchasers of healthcare use aggregated HEDIS® rates to evaluate health insurance companies' efforts to improve preventive health outreach for members.

Physician-specific scores are also used to measure your practice's preventive care efforts. Your practice's HEDIS® score determines your rates for physician incentive programs that pay you an increased premium — for example Pay For Performance or Quality Bonus Funds.



How are rates calculated?

HEDIS® rates can be calculated in two ways: administrative data or hybrid data. Administrative data consists of claim or encounter data submitted to the health plan. Hybrid data consists of both administrative data and a sample of medical record data. Hybrid data requires review of a random sample of member medical records to abstract data for services rendered but that were not reported to the health plan through claims/encounter data. Accurate and timely claim/encounter data reduces the need for medical record review. If services are not billed or not billed accurately, they are not included in the calculation.



How can I improve my HEDIS® scores?

- ✓ Submit claim/encounter data for each and every service rendered
- ✓ Make sure that chart documentation reflects all services billed
- ✓ Bill (or report by encounter submission) for all delivered services, regardless of contract status
- ✓ Ensure that all claim/encounter data is submitted in an accurate and timely manner
- ✓ Include CPT II codes to provide additional details and reduce medical record requests
- ✓ Respond timely to medical records requests
- ✓ Submit supplemental data throughout the measurement year
- ✓ Early Engagement with Pharmacy Adherence is key once a member loses days on a prescription, those days cannot be recovered.
- ✓ Speak with the members about any barriers to adherence.
- ✓ Consider utilizing RxEffect a free online portal for our network providers that will prioritize your high-risk patients more efficiently. This will save on resources as it lists your patients at highest risk for non-adherence.
- ✓ If you have any questions regarding pharmacy and member barriers, please reach out to your local Provider Relations Representative for assistance.





Updates to HEDIS® Measures

(effective for calendar year 2022 and 2023)

This guide has been updated with information from the release of the HEDIS® 2023 Volume 2 Technical Specifications by NCQA and is subject to change.



Retired Measures:

- · (FVA) Flu Vaccinations for Adults Ages 18-64
- (BCS) Breast Cancer Screening: no longer collected administratively; only the BCS-E measure will be reported.
- · (ADV) Annual Dental Visit
- · (FSP) Frequency of Selected Procedures
- (FVO) Flu Vaccinations for Adults Ages 65 and Older
- · (PNU) Pneumococcal Vaccination Status for Older Adults



Revised Measures:

- (FUA) Follow-Up After Emergency Department Visit for Substance Use
- (IET) Initiation and Engagement of Substance Abuse Disorder Treatment
- · (BPD) Blood Pressure Control for Patients with Diabetes
- \cdot (EED) Eye Exam for Patients with Diabetes
- (HBD) Hemoglobin A1c Control for Patients with Diabetes
- \cdot (SPD) Statin Therapy for Patients with Diabetes
- · (AMR) Asthma Medication Ratio
- · (PCE) Pharmacotherapy Management of COPD Exacerbation
- · (BCS-E) Breast Cancer Screening
- · (CCS) Cervical Cancer Screening
- \cdot (PPC) Prenatal and Postpartum Care
- · (PRS-E) Prenatal Immunization Status



New Measure for Medicaid:

· (COL) Colorectal Cancer Screening



Quick Reference Guide

Contents

Adult Health

	Health Services	10
ia	(ACP) Advance Care Planning	.11
SUL A	(AIS) Adult Immunization Status	.11
	(BPD) Blood Pressure Control for Patients with Diabetes	.12
	(CBP) Controlling High Blood Pressure	14
22	(COA) Care for Older Adults	15
ŤŤ	(COL) Colorectal Cancer Screening	16
©	(EED) Eye Exam for Patients with Diabetes	17
0	(FMC) Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions	18
	(HBD) Hemoglobin A1c Control for Patients with Diabetes	20
6/2	(KED) Kidney Health Evaluation for Patients with Diabetes	22
*	(PBH) Persistence of Beta-Blocker Treatment After a Heart Attack	23

3	(PCE) Pharmacotherapy Management of COPD Exacerbation	24
Ų,	(PCR) Plan All Cause Readmission	25
*	(SPC) Statin Therapy for Patients with Cardiovascular Disease	25
	(SPR) Use of Spirometry Testing in the Assessment and Diagnosis of COPD	27
My.	(TRC) Transitions of Care	28
Pha	rmacy Adherence Measures	
	(DIAB) Adherence to Diabetes Medications – Measure Overview	30
	(RASA) Adherence to Hypertensive Medications – Measure Overview	31
	(STAT) Adherence to Cholesterol Medications – Measure Overview	32
	(SUPD) Statin Use in Persons with Diabetes – Measure Overview	33
Wor	nen's Health	
%	(BCS-E) Breast Cancer Screening	. 34
8	(CCS) Cervical Cancer Screening	35
/O \	(CHL) Chlamydia Screening in Women	37
000	(OMW) Osteoporosis Management in Women Who Had a Fracture	38
	(OSW) Osteoporosis Screening in Older Women	39

\$	(PPC) Prenatal and Postpartum Care	10
A SELLA	(PRS-E) Prenatal Immunization Status	41
<u>Pedi</u>	<u>iatric Health</u>	
	(AAB) Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis	42
	(CIS) Childhood Immunization Status	43
	(IMA) Immunizations for Adolescents	45
	(LSC) Lead Screening in Children	45
	(URI) Appropriate Treatment for Upper Respiratory Infection	46
	(W30/WCV) Well-Child and Adolescent Well-Care Visits	46
	(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents	48
<u>Gen</u>	<u>eral Health</u>	
(≥• 1	(AMR) Asthma Medication Ratio	19
	(CWP) Appropriate Testing for Pharyngitis	52
Soci	al Determinants of Health (SDOH)	
	(SNS-E) Social Need Screening and Intervention	53

Behavioral Health

*	(ADD) Follow-up Care for Children Prescribed ADHD Medication	55
	(AMM) Antidepressant Medication Management	
R	(APP) Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics	60
R	(COU) Risk of Continued Opioid Use	60
~	(FUA) Follow-Up After Emergency Department Visit with Substance Use Disorder	61
©	(FUH) Follow-Up After Hospitalization for Mental Illness	65
^	(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder	68
<u></u>	(FUM) Follow-Up After Emergency Department Visit for Mental Illness	72
T/p	(IET) Initiation and Engagement of Substance Use Disorder Treatment	76
®	(POD) Pharmacotherapy for Opioid Use Disorder	79
R	(SAA) Adherence to Antipsychotic Medications for Individuals With Schizophrenia	80
∞	(SSD) Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications	81
0	(UOP) Use of Opioids from Multiple Providers	82

For a complete list of codes, please visit the NCQA website at www.ncqa.org, or see the HEDIS value sets. Only subsets of the NCQA approved codes are listed in this document.

Adult Health

<u>Call To Action</u>: Please refer to the provider portal where you will find a complete list of member care gaps as applicable for the measures in this document.



(AAP) Adults' Access to Preventive/ Ambulatory Health Services

Measure evaluates the percentage of members 20 years and older who had an ambulatory or preventive care visit. Services that count include outpatient evaluation and management (E&M) visits, consultations, assisted living/home care oversight, preventive medicine, and counseling.

- Assist or schedule member's appointments for Preventive Care Visits.
- · Document the date and the type of visit.
- · Submit the applicable codes.

CPT*	HCPCS*	ICD-10*
99202-99205, 99211-99215, 99241- 99245, 99341-99345, 99347- 99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99429, 92002, 92004, 92012, 92014, 99304-99310, 99315, 99316, 99318, 99324-99328, 99334-99337, 98966-98968, 99441-99443, 98969, 98970, 98971, 98972, 99421, 99422, 99423, 99444, 99457, 99483	G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2061, G2062, G2063, T1015, S0620, S0621	Z00.00, Z00.01, Z00.3, Z00.5, Z00.8, Z00.121, Z00.129, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.79, Z02.81, Z02.82, Z02.83, Z02.89, Z02.9, Z76.1, Z76.2

^{*}Codes subject to change





(ACP) Advance Care Planning

Measure evaluates percentage of adults:

- √ 66 and up years of age with advanced illness, an indication of frailty, or who are receiving palliative care and had advance care planning;
- √ 81 years of age and older who had advance care planning

Tips:

- Encourage members to consider an Advance Directive.
- · Assist members in scheduling an Annual Well-visit.
- · Telephone visits, e-visits, or virtual check-ins are acceptable.
- · Submit the applicable codes.

Description	Codes*
Advanced Care Planning	CPT: 99483, 99497
	CPT-CAT-II: 1123F, 1124F, 1157F, 1158F
	HCPCS: S0257
	ICD-10: Z66

^{*}Codes subject to change



(AIS) Adult Immunization Status

Measures percentage of members 19 years of age and older who are up to date on recommended routine vaccines for influenza, tetanus, and diphtheria (Td) or tetanus, diphtheria, and acellular pertussis (Tdap), zoster and pneumococcal.

Tips:

- · Schedule appointments within immunization timeframes.
- Discuss the importance of vaccinations during member appointments.
- Include immunization history from all sources in the member's medical record.



(AIS) Adult Immunization Status (continued)

Description	Codes*
Adult Influenza Vaccine Procedure	CPT: 90630, 90653-90654, 90656, 90658, 90661-90662, 90673-90674, 90682, 90686, 90688-90689, 90694, 90756
Td Vaccine Procedure	CPT: 90714, 90718
Tdap Vaccine Procedure	CPT: 90715
Varicella Zoster (VZV) Vaccine Procedure	CPT: 90710, 90716

^{*}Codes subject to change



(BPD) Blood Pressure Control for Patients with Diabetes

Measure evaluates percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose blood pressure was adequately controlled (<140/90 mm Hg).

Tips:

- · Never round up BP readings.
- · Use correct cuff size on bare arm.
- Check BP on both arms and record the lowest systolic and diastolic readings.
- Patients should rest quietly for at least 5 minutes before the first BP is taken.
- · Submit applicable codes.

Description	Codes*
Palliative Care	HCPCS: G9054, M1017
Outpatient Codes (must include a diagnosis of diabetes)	CPT: 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99987, 99391-99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015



(BPD) Blood Pressure Control for Patients with Diabetes (continued)

Description	Codes*
Non-Acute Inpatient (must include a diagnosis of diabetes)	CPT: 99304-99310, 99315, 99316, 99318, 99324-99328, 99334-99337
Telephone Visits (must include a diagnosis of diabetes)	CPT: 98966-98968, 99441-99443
E-Visits or Virtual Check-ins (must include a diagnosis of diabetes)	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063
Systolic Greater Than/ Equal to 140	CPT-CAT-II: 3077F
Systolic Less Than 140	CPT-CAT-II: 3074F, 3075F
Diastolic 80-89	CPT-CAT-II: 3079F
Diastolic Greater Than/ Equal to 90	CPT-CAT-II: 3080F
Diastolic Less Than 80	CPT-CAT-II: 3078F

^{*}Codes subject to change



(CBP) Controlling High Blood Pressure

Measure evaluates the percentage of members 18-85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (<140/90 mm Hg).

- Blood pressure reading can be collected via any telehealth visit and it does not require a remote monitoring device to be the source.
- Retake BP readings if the reading is >140/90 mm Hg.
- Help members schedule their hypertension follow-up appointments.
- Educate members on what a controlled blood pressure means.
- Talk with members about taking their own blood pressure via a digital device.
- · Submit applicable codes.

Description	Codes*
Essential Hypertension	ICD-10: I10
Systolic Greater Than/Equal to 140	CPT-CAT-II: 3077F
Systolic Less Than 140	CPT-CAT-II: 3074F, 3075F
Diastolic Greater Than/Equal to 90	CPT-CAT-II: 3080F
Diastolic 80-89	CPT-CAT-II: 3079F
Diastolic Less Than 80	CPT-CAT-II: 3078F
Telephone Visits	CPT: 98966-98968, 99441-99443
Palliative Care	HCPCS: G9054, M1017

^{*}Codes subject to change



(COA) Care for Older Adults

Measure evaluates percentage of adults 66 years and older who had each of the following:

✓ Medication review

✓ Functional status assessment

✓ Pain assessment

- A Functional Status Assessment does not require a specific setting. Services rendered during a telephone visit, e-visit, or virtual check-in meets criteria.
- · Submit applicable codes.

Description	Codes*
Medication Review (would need both CPT-CAT II codes to get credit) 1159F (Medication List) & 1160F (Medication Review)	CPT: 90863, 99605, 99606, 99483, 99495, 99496 CPT-CAT-II: 1159F, 1160F HCPCS: G8427
Functional Status Assessment	CPT: 99483 CPT-CAT-II: 1170F HCPCS: G0438, G0439
Pain Assessment	CPT-CAT-II: 1125F, 1126F

^{*}Codes subject to change




(COL) Colorectal Cancer Screening

Measure evaluates the percentage of members 45-75 years of age who has had an appropriate screening for colorectal cancer.

- · Complete and document all screenings for patients.
- Educate members on the importance of colorectal cancer screenings for early detection and the options available to complete their screening.
- Talk with members about using the home screenings for colorectal cancer screening.
- Help members schedule their colonoscopy screening appointments.
- · Submit applicable codes.

Description	Codes*
Colonoscopy	CPT: 44388-44394, 44397, 44401-44408, 45355, 45378-45393, 45398 HCPCS: G0105, G0121
CT Colonography	CPT: 74261-74263
sDNA FIT Lab Test	CPT: 81528
Flexible Sigmoidoscopy	CPT: 45330-45335, 45337-45338, 45340-45342, 45346-45347, 45349-45350 HCPCS: G0104
FOBT Lab Test	CPT: 82270, 82274 HCPCS: G0328
Colorectal Cancer	ICD-10: C18.0-C18.9, C19, C20, C21.2, C21.8, C78.5, Z85.038, Z85.048
Palliative Care	HCPCS: G9054, M1017
Total Colectomy	CPT: 44150-44153, 44155-44158, 44210-44212
*Codes subject to change	



Measure evaluates percentage of members 18-75 years of age with diabetes (type 1 and type 2) who had a retinal eye exam.

- Refer diabetic members to an acceptable eye care professional annually.
- Educate members on the eye damage that could be caused by their diabetes.
- Help members to schedule their annual diabetic eye exam appointments.
- · Submit applicable codes.

Description	Codes*
Palliative Care	HCPCS: G9054, M1017
Outpatient Codes (must include a diagnosis of diabetes)	CPT: 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015
Non-Acute Inpatient (must include a diagnosis of diabetes)	CPT: 99304-99310, 99315, 99316, 99318, 99324-99328, 99334-99337
Telephone Visits (must include a diagnosis of diabetes)	CPT: 98966-98968, 99441-99443
E-Visits or Virtual Check-ins (must include a diagnosis of diabetes)	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063
Unilateral Eye Enucleation With a Bilateral Modifier	CPT: 65091, 65093, 65101, 65103, 65105, 65110,65112, 65114 CPT Modifier: 50 (continued)



(EED) Eye Exam for Patients with Diabetes (continued)

Description	Codes*
Automated Eye Exam	CPT: 92229
Diabetic Retinal Screening Negative in Prior Year	CPT-CAT-II: 3072F
Eye Exam With Retinopathy	CPT-CAT-II: 2022F, 2024F, 2026F HCPCS: S0620, S0621, S3000
Eye Exam Without Retinopathy	CPT-CAT-II: 2023F, 2025F, 2033F

^{*}Codes subject to change



(FMC) Follow-Up After Emergency **Department Visit for People with Multiple High-Risk Chronic Conditions**

The percentage of emergency department (ED) visits for members 18 years of age and older who have multiple high-risk chronic conditions who had a follow-up service within 7 days of the ED visit.

Tips:

- If a patient has more than one ED visit, they could be in the measure more than once.
- Maintain reserved appointments so patients with an ED visit can be seen within 7 days of their discharge.
- · An in-person office visit is not required, follow-up can be provided via a telehealth, telephone, e-visit or virtual visit.
- Submit applicable codes.



(FMC) Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic **Conditions** (continued)

FMC Value Set		
Complex Care Management Services	G0506	
Online Assessments	G0071, G2010, G2012, G2061-G2063, G2250-G2252	
Outpatient	G0402, G0438-G0439, G0463, T1015	
Case Management Encounter	T1016-T1017, T2022-T2023	
Substance Use Disorder Services	G0396-G0397, G0443, H0001, H0005, H0007, H0015-H0016, H0022, H0047, H0050, H2035- H2036, T1006, T1012	
BH Outpatient	G0155, G0176-G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036-H0037, H0039-H0040, H2000, H2010-H2011, H2013-H2020, T1015	

^{*}Codes subject to change



(HBD) Hemoglobin A1c Control for Patients with Diabetes

Measure evaluates percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose HbA1c was at the following levels:

✓ HbA1c control (<8.0%)
✓ HbA1c Poor control (>9.0%)

Tips:

- Document all A1c lab values with dates for diabetic members.
- · Provide education to members regarding the need to monitor and manage their blood sugars (HgA1c).
- · Assist members if needed to schedule lab visits for regular A1c testing to include transportation assistance.
- · Submit applicable codes.

Description	Codes*
Palliative Care	HCPCS: G9054, M1017
Outpatient Codes (must include a diagnosis of diabetes)	CPT: 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015
Non-Acute Inpatient (must include a diagnosis of diabetes)	CPT: 99304-99310, 99315, 99316, 99318, 99324-99328, 99334-99337
Telephone Visits (must include a diagnosis of diabetes)	CPT: 98966-98968, 99441-99443
E-Visits or Virtual Check-ins (must include a diagnosis of diabetes)	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063
HbA1c Lab Test	CPT: 83036, 83037



(HBD) Hemoglobin A1c Control **for Patients with Diabetes** (continued)

Description	Codes*
HbA1c Level Less than 7 Codes	CPT-CAT-II: 3044F
HbA1c Level Greater Than/Equal to 7 and Less than 8	CPT-CAT-II: 3051F
HbA1c Level Greater Than/Equal to 8 and Less Than/Equal to 9	CPT-CAT-II: 3052F
HbA1c Greater Than 9.0	CPT-CAT-II: 3046F

^{*}Codes subject to change



(KED) Kidney Health Evaluation for Patients with Diabetes

The percentage of members 18–85 years of age with diabetes (Type 1 and Type 2) who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) *and* a urine albumin-creatinine ratio (uACR), during the measurement year.

- Use CPT II coding when completing screening test to assist in administrative collection and gap closure.
- Educate members on why good kidney function is important as they work to manage their health and diabetes.
- Help members schedule their diabetes follow-up appointments and remind them of the care gaps that should be covered to include kidney function.
- · Submit applicable codes.

Description	Codes*
Estimated Glomerular Filtration Rate (eGFR) – must be within 4 days or less of the uACR	CPT: 80047, 80048, 80050, 80053, 80069, 82565
Urine Albumin-Creatinine Ratio (uACR) – must be within 4 days or less of the eGFR	CPT: 82043, 82570
Palliative Care	HCPCS: G9054, M1017

^{*}Codes subject to change



(PBH) Persistence of Beta-Blocker Treatment After a Heart Attack

This measure demonstrates the percentage of members 18 years of age and older during the measurement year who were hospitalized and discharged from July 1 of the year prior to June 30 of the measurement year with a diagnosis of AMI and who received persistent beta-blocker treatment for six months after discharge.

	Beta-Blocke	er Medication		
Description	Prescription			
Noncardioselective Beta-blockers	CarvedilolPropranolol	LabetalolTimolol	NadololSotalol	· Pindolol
Cardioselective Beta-blockers	AcebutololAtenolol	BetaxololBisoprolol	MetoproloNebivolol	l
Antihypertensive Combinations	 Atenolol-chlorthalidone Bendroflumethiazide-nadolol Bisoprolol-hydrochlorothiazide Hydrochlorothiazide-metoprolol Hydrochlorothiazide-propranolol 			



(PCE) Pharmacotherapy Management of COPD Exacerbation

Measure evaluates percentage of COPD exacerbations for members 40 years of age and older who had an acute inpatient discharge or ED visit on or between January 1-November 30 and were dispensed appropriate medications.

Two rates are reported:

- Dispensed a systemic **corticosteroid** (or there was evidence of an active prescription) **within 14 days of the event**
- 2 Dispensed a **bronchodilator** (or there was evidence of an active prescription) within **30 days of the event**

Systemic Corticosteroid Medications				
Description	Prescription			
Glucocorticoids	CortisonePrednisoloneMethylprednisolone	_	ocortisone methasone nisone	
	Bronchodilator M	edicati	ons	
Description	Prescription			
Anticholinergic Agents	Aclidinium-bromideIpratropium	· Ume	clidinium opium	
Beta 2-agonists	AlbuterolMetaproterenolIndacaterol		lbuterol ioterol aterol	ArformoterolSalmeterol
Bronchodilator Combinations	 Albuterol-ipratropium Budesonide-formoterol Formoterol-mometasone Glycopyrrolate-indacaterol Umeclidinium-Vilanterol Olodaterol-tiotropium 		FormoteroFluticasoFluticasoFluticaso	ol-aclidinium ol-glycopyrrolate ne-salmeterol ne-vilanterol ne furoate- nium-vilarterol



(PCR) Plan All Cause Readmission

For members 18 years of age and older, the number of acute inpatient and observation stays during the measurement year that were followed by an unplanned acute readmission for any diagnosis within 30 days and the predicted probability of an acute readmission.

*Note: for commercial (Marketplace) and Medicaid, report only members 18-64 years of age.

Description	Codes*
Acute Inpatient	CPT: 99221-99223, 99231-99233, 99238-99239, 99251-99255, 99291
Nonacute Inpatient	CPT: 99304-99310, 99315-99316, 99318, 99324-99328, 99334-99337
Observation	CPT: 33217-99220

^{*}Codes subject to change



(SPC) Statin Therapy for Patients with Cardiovascular Disease

The percentage of males 21-75 years of age and females 40-75 years of age during the measurement year who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and met the following criteria.

The following rates are reported:

- **Received Statin Therapy:** Members who were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement year.
- 2 **Statin Adherence 80%:** Members who remained on a high-intensity or moderate intensity statin medication for at least 80% of the treatment period.



(SPC) Statin Therapy for Patients with Cardiovascular Disease (continued)

Tips:

- Review medication list during each visit with the patient.
- Discuss the importance of medication adherence with the patient.

High- and Moderate-Intensity Statin Medications		
Description	Prescription	Medication Lists
High-intensity Statin Therapy	· Atorvastatin 40-80 mg	Atorvastatin High Intensity Medications List
High-intensity Statin Therapy	• Amlodipine-atorvastatin 40-80 mg	Amlodipine Atorvastatin High Intensity Medications List
High-intensity Statin Therapy	• Rosuvastatin 20-40 mg	Rosuvastatin High Intensity Medications List
High-intensity Statin Therapy	· Simvastatin 80 mg	Simvastatin High Intensity Medications List
High-intensity Statin Therapy	• Ezetimibe-simvastatin 80 mg	Ezetimibe Simvastatin High Intensity Medications List
Moderate-intensity Statin Therapy	· Atorvastatin 10-20 mg	Atorvastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	• Amlodipine-atorvastatin 10-20 mg	Amlodipine Atorvastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	· Rosuvastatin 5-10 mg	Rosuvastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	· Simvastatin 20-40 mg	Simvastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	• Ezetimibe-simvastatin 20-40 mg	Ezetimibe Simvastatin Moderate Intensity Medication List



(SPC) Statin Therapy for Patients with Cardiovascular Disease (continued)

High- and Moderate-Intensity Statin Medications		
Description	Prescription	Medication Lists
Moderate-intensity Statin Therapy	• Pravastatin 40-80 mg	Pravastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	· Lovastatin 40 mg	Lovastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	• Fluvastatin 40-80 mg	Fluvastatin Moderate Intensity Medications List
Moderate-intensity Statin Therapy	· Pitavastatin 1-4 mg	Pitavastatin Moderate Intensity Medications List



(SPR) Use of Spirometry Testing in the Assessment and Diagnosis of COPD

Measure evaluates the percentage of members 40 years of age and older with a new diagnosis of COPD or newly active COPD, who received appropriate spirometry testing to confirm diagnosis.

CPT*

Spirometry 94010, 94014-94016, 94060, 94070, 94375

^{*}Codes subject to change



(TRC) Transitions of Care

The percentage of discharges for members 18 years of age and older who had each of the following:

✓ Notification of Inpatient Admission.

Documentation of receipt of notification of inpatient admission on the day of admission through 2 days after the admission (3 total days).

✓ Receipt of Discharge Information.

Documentation of receipt of discharge information on the day of discharge through 2 days after the discharge (3 total days).

✓ Patient Engagement After Inpatient Discharge.

Documentation of patient engagement (e.g., office visits, visits to the home, telehealth) provided within 30 days after discharge.

✓ Medication Reconciliation Post-Discharge.

Documentation of medication reconciliation on the date of discharge through 30 days after discharge (31 total days).

- Ensure follow-up appointments are scheduled within 30 days after discharge.
- Use CPT II coding when completing screening test to assist in administrative collection and gap closure.
- · Document medication reconciliation.
- Services may be performed during a telephone visit, e-visit, or virtual check-in.
- · Submit applicable codes.



(TRC) Transitions of Care (continued)

Description	Codes*
Medication Reconciliation	CPT-CAT-II: 1111F CPT: 99483, 99495-99496
Telephone Visits Outpatient	CPT: 99441-99442, 98966-98968 CPT: 99201-99205, 99211-99215, 99241-99245, 99341-99350, 99381-99387, 99391-99397, 99401, 99483
Online Assessments	HCPCS: G8427, G0071, G2010, G2012, G2061-G2063, G2250-G2252
Transitional Care Management Services	99495, 99496

^{*}Codes subject to change

Pharmacy Adherence Measures



(DIAB) Adherence to Diabetes Medications – Measure Overview

The percentage of members 18 years and older with a diabetes medication with a Proportion of Days Covered (PDC) ≥ 80%.

- ✓ Higher rate indicates better performance
- ✓ 2 fills needed to index into the measure
- ✓ Targeted early in the year

RY25 Measure Weight: 3

Gap Closure Requirements

PDC ≥ 80% per member

- ✓ PDC calculated utilizing: total days supplied of diabetes pharmacy claims / Date of first diabetes fill to the end of the reporting interval
- ✓ Each medication claim must be submitted to the Health Plan (cash payment/ samples given by prescriber or meds filled at an out of network pharmacy do not count)
- ✓ Final plan star score based upon the percentage of members with a PDC ≥ 80%

- ✓ Medication Inclusions: Diabetes Medications
 - · i.e., Metformin, Glipizide, Glimepiride, Januvia
- ✓ Exclusions: Members with an insulin claim; Hospice enrollees, ESRD



(RASA) Adherence to Hypertensive Medications – Measure Overview

The percentage of members 18 years and older with a RASA medication with a Proportion of Days Covered (PDC) \geq 80%.

- ✓ Higher rate indicates better performance
- ✓ 2 fills needed to index into the measure
- ✓ Targeted early in the year

RY25 Measure Weight: 3

Gap Closure Requirements

PDC ≥ 80% per member

- ✓ PDC calculated utilizing: total days supplied of RASA pharmacy claims / Date of first RASA fill to the end of the reporting interval
- ✓ Each medication claim must be submitted to the Health Plan (cash payment/ samples given by prescriber or meds filled at an out of network pharmacy do not count)
- ✓ Final plan star score based upon the percentage of members with a PDC ≥ 80%

- ✓ Medication Inclusions: RASA Medications.
 - · i.e. Lisinopril, Losartan, Enalapril, Valsartan
- ✓ Exclusions: Members with a Sacubutril/valsartan claim; Hospice enrollees, ESRD

(STAT) Adherence to Cholesterol Medications - Measure Overview

The percentage of members 18 years and older with a CHOL medication with a Proportion of Days Covered (PDC) \geq 80%.

- ✓ Higher rate indicates better performance
- ✓ 2 fills needed to index into the measure
- ✓ Targeted early in the year

RY25 Measure Weight: 3

Gap Closure Requirements

PDC ≥ 80% per member

- ✓ PDC calculated utilizing: total days supplied of CHOL pharmacy claims / Date of first CHOL fill to the end of the reporting interval
- ✓ Each medication claim must be submitted to the Health Plan (cash payment/ samples given by prescriber or meds filled at an out of network pharmacy do not count)
- ✓ Final plan star score based upon the percentage of members with a PDC > 80%

- ✓ Medication Inclusions: CHOL Medications
 - · i.e., Atorvastatin, Simvastatin, Rosuvastatin, Pravastatin
- ✓ Exclusions: Hospice enrollees, ESRD

(SUPD) Statin Use in Persons with Diabetes – Measure Overview

The percentage of members ages 40-75 years of age with diabetes that have a single fill of a statin.

- ✓ Higher rate indicates better performance
- ✓ Only 1 fill needed to index in the measure
- ✓ Targeted later in the year vs. other measures (starting in late July or August)

RY25 Measure Weight: 1

Gap Closure Requirements

Member received a Statin Therapy

✓ The number of members who had at least one dispensing event for a statin medication during the measurement year

- ✓ Medication Inclusions: Statin Medications
 - · i.e., Atorvastatin, Simvastatin, Rosuvastatin, Pravastatin
- ✓ Exclusions: ESRD, Rhabdomyolysis, Pregnancy, Cirrhosis, Pre-Diabetes, Polycystic Ovary Syndrome

Women's Health



(BCS-E) Breast Cancer Screening

The Breast Cancer Screening measure has transitioned to exclusive use of the Electronic Clinical Data Systems reporting standard for measurement year 2023.

Summary of Changes:

Only BCS-E measure will be reported. BCS is a retired measure and replaced with the new BCS-E measure.

Measure evaluates the percentage of women 50-74 years of age who had a mammogram to screen for breast cancer.

Tips:

- · Schedule member's mammogram screening.
- Document the date and the specific procedure completed when reviewing the patient's history.
- · Submit applicable codes.



(BCS-E) Breast Cancer Screening (continued)

Description	Codes*
Mammogram	CPT: 77061-77063, 77065-77067
	ICD-10 (bilateral mastectomy): Z90.13
	SNOMED:
	836381000000102
	1106021000000101
	1106641000000102
	1106651000000104
	1106661000000101
	1111381000000105
	1111411000000107
	1111421000000101
	111179100000108
Palliative Care	HCPCS: G9054, M1017

^{*}Codes subject to change



(CCS) Cervical Cancer Screening

This measure demonstrates the percentage of women 21-64 years of age who were screened for cervical cancer using **either** of the following criteria:

- ✓ Women 21-64 years of age who had cervical cytology performed within last 3 years.
- ✓ Women 30-64 years of age who had cervical high-risk human papillomavirus (hrHPV) testing performed within the last 5 years.
- ✓ Women 30-64 years of age who had cervical cytology/high risk human papillomavirus (hrHPV) co-testing within the last 5 years.



(CCS) Cervical Cancer Screening (continued)

- · Help members schedule their routine cervical cancer screening.
- Document the date and the specific procedure completed when reviewing the patient's history.
- · Submit the applicable codes

Description	Codes*
Cervical Cytology Lab Test (20-64)	CPT: 88141-88143, 88147, 88148, 88150, 88152-88153, 88164-88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091
HPV Tests (30-64)	CPT: 87624, 87625 HCPCS: G0476
Hysterectomy with No Residual Cervix and Absence of Cervix Diagnosis	CPT: 57530, 57531, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290-58294, 58548, 58550, 58552-58554, 58570-58573, 58575, 58951, 58953, 58954, 58956, 59135 ICD-10: Q51.5, Z90.710, Z90.712 SNOMED: 1163275000
Palliative Care	HCPCS: G9054, M1017

^{*}Codes subject to change



(CHL) Chlamydia Screening in Women

Measure evaluates the percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia.

Tips:

- · Perform chlamydia screening every year.
- Inform patient that chlamydia screening can be performed through a urine test. Offer this as an option for patients.
- Add chlamydia screening as a standard lab for women 16–24 years old. Use well-child exams and well-women exams for this purpose.
- Place chlamydia swab next to Pap test or pregnancy detection materials.
- Meet with teens and young adults separately from their parents to allow open conversation.
- Advise members during wellness visits or when they are seen for birth control to get screened for chlamydia.
- · Submit applicable codes.

CPT*

87110, 87270, 87320, 87490-87492, 87810, 0353U

*Codes subject to change



(OMW) Osteoporosis Management in Women Who Had a Fracture

Measure evaluates the percentage of women 67-85 years of age who suffered a fracture and who had either a bone mineral density (BMD) test or prescription for a drug to treat osteoporosis in the six months after the fracture.

Tips:

- Provide patients who have had a fracture with a referral for BMD testing and encourage them to obtain the screening.
- When appropriate, prescribe medication to treat osteoporosis (bisphosphates).
- · Check that fracture codes are used appropriately.
- Consider offering onsite bone density screening for patients at risk.
- Women at risk for osteoporosis should receive a bone density screening every two years.
- · Submit applicable codes.

Description	Codes*
Palliative Care	HCPCS: G9054, M1017
Bone Mineral Density Tests	CPT: 76977, 77078, 77080, 77081, 77085, 77086
Osteoporosis Medications	HCPCS: J0897, J1740, J3110, J3111, J3489
Long-Acting Osteoporosis Medications during Inpatient Stay	HCPCS: J0897, J1740, J3489

^{*}Codes subject to change



(OMW) Osteoporosis Management in Women Who Had a Fracture (continued)

Osteoporosis Medications		
Description	Prescription	
Bisphosphonates		Alendronate-cholecalciferolRisedronateZoledronic acid
Other agents	AbaloparatideTeriparatide	RomosozumabRaloxifene



(OSW) Osteoporosis Screening in Older Women

The percentage of women 65-75 years of age who received osteoporosis screening.

Description	Codes*
Osteoporosis Screening Tests	CPT: 76977, 77078, 77080, 77081, 77085
Palliative Care	HCPCS: G9054, M1017

^{*}Codes subject to change



(PPC) Prenatal and Postpartum Care

Measure evaluates percentage of deliveries of live births on or between October 8 of the year prior to the measurement year and October 7 of the measurement year. For these women, the measure assesses the following facets of prenatal and postpartum care.

Tips:

- Schedule an initial prenatal visit within the first 12 weeks of pregnancy with an OB/GYN, PCP, or nurse midwife.
- Educate members on the importance of prenatal care throughout their pregnancy to include the postpartum visit.
- · Schedule postpartum visits prior to discharge after delivery.
- · Submit applicable codes.
- ✓ **Timeliness of Prenatal Care:** percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date, or within 42 days of enrollment in the organization
- ✓ Postpartum Care: percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery

Description	Codes*
Online Assessments	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061, G2062, G2063
Prenatal Visits	CPT: 99201-99205, 99211-99215, 99241-99245, 99483 HCPCS: GO463, T1015
Stand-Alone Prenatal Visits	CPT: 99500 CPT-CAT-II: 0500F, 0501F, 0502F HCPCS: H1000, H1001, H1002, H1003, H1004
Cervical Cytology Lab Test	CPT: 88141-88143, 88147, 88148, 88150, 88152-88153, 88164-88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091



(PPC) Prenatal and Postpartum Care (continued)

Description	Codes*
Postpartum Visits	CPT: 57170, 58300, 59430, 99501
	CPT-CAT-II: 0503F
	HCPCS: G0101
	ICD-10: Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2
Telephone Visits	CPT: 98966-98968, 99441-99443

^{*}Codes subject to change

NOTE: When using the Online Assessment, Telephone Visit, or Prenatal Visit codes, remember to also include a Pregnancy Diagnosis code.



(PRS-E) Prenatal Immunization Status

Measures percentage of women who received influenza, tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccinations in the measurement period.

For a complete list of approved codes, please visit the NCQA website at **www.ncqa.org**.

Description	Codes*
Adult Influenza Immunization	CVX: 88, 135, 140-141, 144, 150, 153, 155, 158, 166, 168, 171, 185-186, 197, 205
Adult Influenza Vaccine Procedure	CPT: 90630, 90653-90654, 90656, 90658, 90661-90662, 90673-90674, 90682, 90686, 90688-90689, 90694, 90756
Tdap Immunization	CVX: 115
Tdap Vaccine Procedure	CPT: 90715

^{*}Codes subject to change

Pediatric Health



(AAB) Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis

Percentage of episodes for members 3 months of age and older with a diagnosis of acute bronchitis/bronchiolitis that did not result in an antibiotic dispensing event.

Tips:

- A higher rate indicates appropriate treatment (percent of episodes that were **not** prescribed an antibiotic).
- If patient needs prescription for antibiotic, include appropriate diagnosis to support use of antibiotic prescribed.
- · Submit applicable codes.

Description	Codes*
Acute Bronchitis	J20.3-J20.9, J21.0, J21.1, J21.8, J21.9
Pharyngitis	J02.0, J02.8, J02.9, J03.00, J03.01, J03.80, J03.81, J03.90, J03.91

^{*}Codes subject to change



(CIS) Childhood Immunization Status

This measure demonstrates the percentage of children 2 years of age who completed immunizations on or before the child's second birthday.

Tips:

- Document both the name of the vaccine and the date it was administered in the medical record.
- · Submit applicable codes.

DTaP (4 dose)	CPT: 90697, 90698, 90700, 90723 CVX: 20, 50, 106, 107, 110, 120, 146
HIB (3 dose)	CPT: 90644, 90647, 90648, 90697, 90698, 90748 CVX: 17, 46, 47, 48, 49, 50, 51, 120, 146, 148
Newborn Hep B (3 dose)	CPT: 90697, 90723, 90740, 90744, 90747, 90748 CVX: 08, 44, 45, 51, 110, 146 HCPCS: G0010 ICD-10: B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11
IPV (3 dose)	CPT: 90697, 90698, 90713, 90723 CVX: 10, 89, 110, 120, 146
MMR (1 dose)	CPT: 90707, 90710 CVX: 03, 94 ICD-10: B05.0, B05.1, B05.2, B05.3, B05.4, B05.81, B05.89, B05.9, B26.0, B26.1, B26.2, B26.3, B26.81, B26.82. B26.83, B26.84, B26.85, B26.89, B26.9, B06.00, B06.01, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9
Pneumococcal Conjugate PCV (4 dose)	CPT: 90670, 90671 CVX: 109, 133, 152 HCPCS: G0009



(CIS) Childhood Immunization Status (continued)

Description	Codes*
Varicella VZV (1 dose)	CPT: 90710, 90716 CVX: 21, 94 ICD-10: B01.0, B01.11, B01.12, B01.2, B01.81, B01.89, B01.9, B02.0, B02.1, B02.21, B02.22, B02.23, B02.24, B02.29, B02.30, B02.31, B02.32, B02.33, B02.34, B02.39, B02.7, B02.8, B02.9
Hep A (1 dose)	CPT: 90633 CVX: 31, 83, 85 ICD-10: B15.0, B15.9
Influenza Flu (2 dose) LAIV vaccination must be administered on the child's 2nd birthday	CPT: 90655, 90657, 90660, 90661, 90672, 90673, 90674, 90685-90689, 90756 CVX: 88, 140, 141, 150, 153, 155, 158, 161, 111, 149, 171, 186 HCPCS: G0008
Rotavirus (2 Dose)	CPT: 90681 CVX: 119
Rotavirus (3 Dose)	CPT: 90680 CVX: 116, 122
Anaphylaxis	Use applicable SNOMED as indicated per vaccine

^{*}Codes subject to change

NOTE: Rotavirus is either 2 dose **OR** 3 dose for compliancy



(IMA) Immunizations for Adolescents

Tips:

- Document both the name of the vaccine and the date it was administered in the medical record.
- · Submit applicable codes.

Measure evaluates percentage of adolescents 13 years of age who completed immunizations on or before the member's 13th birthday.

Description	Codes*
Meningococcal – serogroup A,C,W, and Y: (1 dose)	CPT: 90619, 90733, 90734 CVX: 32, 108, 114, 136, 147, 167, 203
Tdap (1 dose)	CPT: 90715 CVX: 115
HPV (2 or 3 dose series)	CPT: 90649-90651 CVX: 62, 118, 137, 165
Anaphylaxis	Use applicable SNOMED as indicated per vaccine
*Codes subject to change	



(LSC) Lead Screening in Children

Measure evaluates percentage of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning on or prior to their second birthday (two tests by age 2).

Tips:

- $\cdot\,$ LSC testing on or prior to the child's second birthday.
- · Document both the date and results of the LSC screening.
- Any missed tests must be completed as soon as possible, but no later than six years of age.
- · Submit applicable codes.

CPT*

83655

*Codes subject to change





(URI) Appropriate Treatment for Upper Respiratory Infection

The percentage of episodes for members 3 months of age and older with a diagnosis of upper respiratory infection (URI) that did not result in an antibiotic dispensing event.

Tips:

- Discourage the use of antibiotics for routine treatment of uncomplicated acute bronchitis, unless clinically indicated.
- · Submit applicable codes.

Description	Codes*
Acute Nasopharyngitis (common cold)	ICD-10: J00
Acute Laryngopharyngitis	ICD-10: J06.0
Acute Upper Respiratory Infection, unspecified	ICD-10: J06.9

^{*}Codes subject to change



(W30/WCV) Well-Child and Adolescent Well-Care Visits

One or more comprehensive well-care visits with a PCP or OB/GYN within the measurement year. Visits occurring anytime in the measurement year, including prior to or after the patient's birthday, closes the gap.

(W30) Well-Child Visits in the First 30 Months of Life:

Children who had the following number of well-child visits with PCP during the last 15 months.



(W30/WCV) Well-Child and Adolescent Well-Care Visits (continued)

Tips:

- Remind caregivers of appointments by texts or phone calls.
- Educate the caregiver about the importance of preventive care visits.
- Consider using templates with checkboxes to ensure required information is documented.
- · Submit applicable codes.

The following rates are reported:

- 1 Well-Child Visits in the First 15 Months. Children who turned 15 months old during the measurement year: Six or more well-child visits.
- 2 Well-Child Visits for Age 15 Months-30 Months. Children who turned 30 months old during the measurement year: Two or more well-child visits.

CPT*	HCPCS*	ICD 10*
99381, 99382, 99391,	G0438, G0439,	Z00.110, Z00.111, Z00.121,
99392, 99461	S0302	Z00.129, Z00.2, Z76.1, Z76.2

^{*}Codes subject to change

(WCV) Child and Adolescent Well-Care Visits: Members 3-21 years of age who had a least one comprehensive well-care visit with a PCP or an OB/GYN

CPT*	HCPCS*	ICD 10*
99382-99385,	G0438, G0439,	Z00.00, Z00.01, Z00.121, Z00.129,
99391-99395	S0302, S0610,	Z00.2, Z00.3, Z01.411, Z01.419,
	S0612, S0613	Z02.5, Z76.2

^{*}Codes subject to change



(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents

This measure demonstrates the percentage of members 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of the following:

- ✓ BMI Percentile
- ✓ Counseling for Nutrition
- ✓ Counseling for physical activity

Tips:

- Be sure to document all components of the WCC measure on every visit.
- · Call members/caregivers to reschedule cancelled appointments.
- · Submit applicable codes.

Description	Codes*
BMI Percentile	ICD-10: Z68.51, Z68.52, Z68.53, Z68.54
Nutrition Counseling	CPT: 97802-97804 HCPCS: G0270, G0271, G0447, S9449, S9452, S9470 ICD-10: Z71.3
Physical Activity	HCPCS: G0447, S9451 ICD-10: Z02.5, Z71.82

^{*}Codes subject to change

General Health



(AMR) Asthma Medication Ratio

Measure evaluates the percentage of members 5-64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medication of 0.50 or greater.

- **Step 1:** For each member, count the units of asthma controller medications (Asthma Controller Medications List) dispensed during the measurement year.
- **Step 2:** For each member, count the units of asthma reliever medications (Asthma Reliever Medications List) dispensed during the measurement year.
 - ✓ For each member, sum the units calculated in step 1 and step 2 to determine units of total asthma medications
 - ✓ For each member, calculate ratio using the below:
 - Units of Controller Medications/Units of Total Asthma Medications

Asthma Controller Medications			
Description	Prescriptions	Medication Lists	Route
Antibody Inhibitors	· Omalizumab	Omalizumab Medications List	Injection
Anti-interleukin-4	· Dupilumab	Dupilumab Medications List	Injection



(AMR) Asthma Medication Ratio (continued)

	Asthma Controller	Medications	
Description	Prescriptions	Medication Lists	Route
Anti-interleukin-5	• Benralizumab	Benralizumab Medications List	Injection
Anti-interleukin-5	· Mepolizumab	Mepolizumab Medications List	Injection
Anti-interleukin-5	· Reslizumab	Reslizumab Medications List	Injection
Inhaled Steroid Combinations	 Budesonide- formoterol 	Budesonide Formoterol Medications List	Inhalation
Inhaled Steroid Combinations	· Fluticasone- salmeterol	Fluticasone Salmeterol Medications List	Inhalation
Inhaled Steroid Combinations	· Fluticasone- vilanterol	Fluticasone Vilanterol Medications List	Inhalation
Inhaled Steroid Combinations	· Formoterol- mometasone	Formoterol Mometasone Medications List	Inhalation
Inhaled Corticosteroids	Beclomethasone	Beclomethasone Medications List	Inhalation
Inhaled Corticosteroids	· Budesonide	Budesonide Medications List	Inhalation
Inhaled Corticosteroids	· Ciclesonide	Ciclesonide Medications List	Inhalation
Inhaled Corticosteroids	• Flunisolide	Flunisolide Medications List	Inhalation
Inhaled Corticosteroids	· Fluticasone	Fluticasone Medications List	Inhalation



(AMR) Asthma Medication Ratio (continued)

	Asthma Controlle	r Medications	
Description	Prescriptions	Medication Lists	Route
Inhaled Corticosteroids	Mometasone	Mometasone Medications List	Inhalation
Leukotriene Modifiers	• Montelukast	Montelukast Medications List	Oral
Leukotriene Modifiers	· Zafirlukast	Zafirlukast Medications List	Oral
Leukotriene Modifiers	· Zileuton	Zileuton Medications List	Oral
Methylxanthines	• Theophylline	Theophylline Medications List	Oral
	Asthma Reliever	Medications	
Description	Prescriptions	Medication Lists	Route
Short-acting, Inhaled Beta-2 Agonists	Albuterol	Albuterol Medications List	Inhalation
Short-acting, Inhaled Beta-2 Agonists	Levalbuterol	Levalbuterol Medications List	Inhalation

(CWP) Appropriate Testing for Pharyngitis

This measure demonstrates the percentage of episodes for members 3 years and older where the member was diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode.

Tips:

- Review and document the group A streptococcus (strep) test in the member's health record.
- An in-person office visit is not required, follow-up can be provided via a telehealth, telephone, e-visit or virtual visit.
- Provide tips for managing viral infections and their symptoms such as over the counter medications.
- · Submit applicable codes.

CPT*

87070, 87071, 87081, 87430, 87650-87652, 87880

*Codes subject to change

Social Determinants of Health (SDOH)



(SNS-E) Social Need Screening and Intervention

This measure demonstrates the percentage of members who were screened, using prespecified instruments, at least once during the measurement period for unmet food, housing and transportation needs, and received a corresponding intervention if they screened positive.

Description	Codes*
Occupational Exposure to Risk Factors	ICD-10: Z57.0 - Z57.9
Problems Related to Education and Literacy	ICD-10: Z55.0 – Z55.9
Problems Related to Employment and Unemployment	ICD-10: Z56.0 – Z56.9
Problems Related to Physical Environment	ICD-10: Z58.0 - Z58.9
Problems Related to Housing and Economic Circumstances	ICD-10: Z59.0 – Z59.9
Problems Related to Social Environment	ICD-10: Z60.0 - Z60.9
Problems Related to Upbringing	ICD-10: Z62.0 - Z62.9
Problems Related to Primary Support Group, Including Family Circumstances	ICD-10: Z63.0 – Z63.9
Problems Related to Certain Psychosocial Circumstances	ICD-10: Z64.0 – Z64.4
(Sport) Special Determinants of Sychosocial Circumstances	of Health (continued)
Problems Related to Substance Use	ICD-10: Z71.41, Z71.42, Z71.51, Z71.52
Problems Related to Sleep/Sleep Hygiene	ICD-10: Z72.820, Z72.821

(SNS-E) Social Need Screening and Intervention (continued)

Description	Codes*
Other Risk Factors	ICD-10: Z91.89
Patient/Caregiver Noncompliance with Dietary Regimen or Medical Treatment Due to Financial Hardship	ICD-10: Z911.10, Z911.90, Z91A.10, Z91A.20
Transportation Insecurity Procedures	CPT: 96156
CPT/HCPCS Screening Codes Applicable to SDOH	CPT: 96156-96161, 97802-97804, 99377-99378 HCPCS : S5170, S9470, G0182, G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046

Best Practices Include supplemental codes in the patient's diagnosis section on a claim form. Assign as many SDOH codes necessary to describe all the social problems, conditions, or risk factors documented during the current episode of care.

^{*}Codes subject to change

Behavioral Health



(ADD) Follow-up Care for Children Prescribed ADHD Medication

Measure evaluates percentage of children newly prescribed attention deficit hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.

Tips:

- Complete a medical and psychiatric examination prior to diagnosing and prescribing ADHD medications.
- Have member or parent schedule a follow-up appointment before leaving the office when a new ADHD medication has been prescribed.
- Have staff reschedule canceled appointments.
- · Schedule telehealth visits if office visits are not acceptable.
- · Submit applicable codes.

Two rates are reported:

- 1 Initiation Phase: percentage of members 6-12 years of age with a prescription dispensed for ADHD medication, who had one follow-up visit with a practitioner with prescribing authority during the 30-day Initiation Phase
- 2 Continuation and Maintenance (C&M) Phase: percentage of members 6-12 years of age with a prescription dispensed for ADHD medication, who remained on the medication for at



(ADD) Follow-up Care for Children Prescribed ADHD Medication (continued)

least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended

Description	Codes*
An Outpatient Visit	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
BH Outpatient Visit	CPT: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99510, 99483, 99492-99494 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015
Observation Visit	CPT: 99217-99220
Health and Behavior Assessment/ Intervention	CPT: 96150-96154, 96156, 96158, 96159, 96164, 96165, 96167, 96168, 96170, 96171
Visit Setting Unspecified Value Set with Partial Hospitalization POS	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 52
Partial Hospitalization/ Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485



(ADD) Follow-up Care for Children Prescribed ADHD Medication (continued)

Description	Codes*
Telehealth Visit	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 02, 10
Telephone Visits	CPT: 98966-98968, 99441-99443
E-visit/Virtual Check-In	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061, G2062, G2063
Visit Setting Unspecified Value Set with Community Mental Health Center POS	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 53
Narcolepsy	ICD-10: G47.411, G47.419, G47.421, G47.429

^{*}Codes subject to change



(AMM) Antidepressant Medication Management

Applicable Foster Care Measure:

Measure evaluates percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication treatment.

Two rates are reported:

- **Effective Acute Phase Treatment:** percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks)
- 2 Effective Continuation Phase Treatment: percentage of members who remained on an antidepressant medication for at least 180 days (6 months)



(AMM) Antidepressant Medication Management (continued)

Antidepressant Medications			
Description	Prescription		
Miscellaneous Antidepressants	BupropionVortioxetine	· Vilazodone	
Monoamine Oxidase Inhibitors	IsocarboxazidPhenelzine	SelegilineTranylcypromine	
Phenylpiperazine Antidepressants	· Nefazodone	· Trazodone	
Psychotherapeutic Combinations	Amitriptyline-chlordiazepoxideFluoxetine-olanzapineAmitriptyline-perphenazine		
SNRI Antidepressants	DesvenlafaxineVenlafaxine	DuloxetineLevomilnacipran	
SSRI Antidepressants	CitalopramFluvoxamine	EscitalopramParoxetine	FluoxetineSertraline
Tetracyclic Antidepressants	• Maprotiline	• Mirtazapine	
Tricyclic Antidepressants	AmitriptylineDesipramineNortriptyline	AmoxapineDoxepin (>6 mg)Protriptyline	ClomipramineImipramineTrimipramine



(APM) Metabolic Monitoring for Children and Adolescents on Antipsychotics

Applicable Foster Care Measure:

This measure demonstrates the percentage of children and adolescents 1-17 years of age who had two or more antipsychotic prescriptions and had metabolic testing.



(APM) Metabolic Monitoring for Children and Adolescents on Antipsychotics (continued)

Tips:

- Provide members/caregivers with lab orders for HbA1c or glucose and cholesterol or LDL-C to be completed yearly.
- Educate the member and caregiver about the risks associated with taking antipsychotic medications and the importance of regular follow up care.
- · Submit applicable codes.

Three rates reported:

- Percentage of children and adolescents on antipsychotics who received blood glucose testing
- Percentage of children and adolescents on antipsychotics who received cholesterol testing
- 3 Percentage of children and adolescents on antipsychotics who received blood glucose and cholesterol testing

Description (Need either A1c or Glucose and LCL C or Cholesterol)	Codes*
HbA1c Lab Tests	CPT: 83036, 83037 CPT-CAT-II: 3044F, 3046F, 3051F, 3052F
Glucose Lab Tests	CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951
LDL-C Lab Tests	CPT: 80061, 83700, 83701, 83704, 83721 CPT-CAT-II: 3048F, 3049F, 3050F
Cholesterol Lab Tests	CPT: 82465, 83718, 83722, 84478

^{*}Codes subject to change



(APP) Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics*

The percentage of children and adolescents 1–17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment.

СРТ	HCPCS
90832 - 90834, 90836 - 90840,	G0176, G0177, G0409 - G0411, H0004,
90845 - 90847, 90849, 90853,	H0035 - H0040, H2000, H2001,
90875, 90876, 90880	H2011 - H2014, H2017 - H2020, S0201,
	S9480, S9484, S9485 With or without
	a telehealth modifier: 95, GT

^{*}Codes subject to change



(COU) Risk of Continued Opioid Use

Measure evaluates the percentage of members 18 years of age and older who have a new episode of opioid use that puts them at risk for continued opioid use.

Two rates are reported:

- 1 The percentage of members with at least 15 days of prescription opioids in a 30-day period.
- 2 The percentage of members with at least 31 days of prescription opioids in a 62-day period.

Note: A lower rate indicates better performance.

Opioid Medications		
 Benzhydrocodone 	 Fentanyl 	 Morphine
· Buprenorphine (transdermal	 Hydrocodone 	Opium
patch and buccal film)	 Hydromorphone 	 Oxycodone
 Butorphanol 	 Levorphanol 	 Pentazocine
· Codeine	 Meperidine 	 Tapentadol
 Dihvdrocodeine 	 Methadone 	 Tramadol



(FUA) Follow-Up After Emergency Department Visit with Substance Use Disorder

Measure evaluates the percentage of emergency department (ED) visits among members age 13 years and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose.

Two rates are reported:

- 1 Discharges for which the member received follow-up within 30 days of discharge
- 2 Discharges for which the member received follow-up within 7 days of discharge

The visit can be with any practitioner if the claim includes a diagnosis of SUD (F10.xx-F19.xx) or drug overdose (e.g.T40-T43, T51). If the visit occurs with a mental health provider, the claim does not have to include the SUD or drug overdose diagnosis.

Description

Outpatient Visit with any Diagnosis of SUD or Drug Overdose

Codes*

CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211- 99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 994883, 99492-99494, 99510 **HCPCS:** G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx

POS: 03, 05, 07, 09, 11-20, 22, 33, 49,

50, 71-72



(FUA) Follow-Up After Emergency Department Visit with Substance Use Disorder (continued)

Intensive Outpatient Encounter or Partial Hospitalization with any Diagnosis of SUD or Drug Overdose **CPT:** 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255

HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485

ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx

POS: 52

Non-residential Substance Abuse Treatment Facility with any Diagnosis of SUD or Drug Overdose **CPT:** 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239,

99251-99255

ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx

POS: 57. 58

Community Mental Health Center Visit with any Diagnosis of SUD or Drug Overdose **CPT:** 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239,

99251-99255

ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx

POS: 53

Observation Visit with any Diagnosis of SUD or Drug Overdose **CPT:** 99217, 99218, 99219, 99220 **ICD-10:** F10.xx-F19.xx or T40. xxxx-T43.xxxx. T51.xxxx



(FUA) Follow-Up After Emergency Department Visit with Substance Use Disorder (continued)

Description	Codes*
Peer Support Service with any Diagnosis of SUD or Drug Overdose	HCPCS: G0177, H0024, H0025, H0038-H0040, H0046, H2014, H2023, S9445, T1012, T1016 ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx
Opioid Treatment Service That Bills Monthly or Weekly with any Diagnosis of SUD or Drug Overdose	HCPCS: G2086, G2087, G2071, G8074-G2077, G2080 ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx
Telehealth Visit with any Diagnosis of SUD or Drug Overdose	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx POS: : 02, 10
Telephone Visit with any Diagnosis of SUD or Drug Overdose	CPT: 98966-98968, 99441-99443 ICD-10: F10.xx-F19.xx or T40. xxxx-T43.xxxx, T51.xxxx
E-Visit or Virtual Check In with any Diagnosis of SUD or Drug Overdose	CPT: 98969-98972, 99421-99444, 99457, 99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252
Substance Use and Substance Use Disorder Services	CPT: 99408, 99409 HCPCS: T1012, G0396, G0397, H0001, H0005, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, H0006, H0028



(FUA) Follow-Up After Emergency Department Visit with Substance Use Disorder (continued)

Description	Codes*
Behavioral Health Screening or Assessment for SUD or Mental Health Disorders	CPT: 99408, 99409 HCPCS: G0396, G0397, G0442, H2011, H0001, H0002, H0031, H0049
Pharmacotherapy Dispensing Event or Medication Treatment Event	Medications: Disulfiram (oral), Naltrexone (oral and injectable), Acamprosate (oral; delayed-release tablet), Buprenorphine (implant, injection, or sublingual tablet), Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) HCPCS: G2069, G2070, G2072, G2073, H0020, H0033, J0570-J0575, J2315, Q9991, Q9992, S0109
*Codes subject to change	



(FUH) Follow-Up After Hospitalization for Mental Illness

Applicable Foster Care Measure:

Measure evaluates percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider.

Tips:

- Schedule follow up appointments prior to discharge and include the date and time on discharge instructions.
- · Submit applicable codes.

	FUH Age Stratification
	Eligible Population
Product Lines:	Commercial (Marketplace), Medicaid, Medicare (report each product line separately)
Ages:	 6 years and older as of the date of discharge. Report three ag stratifications and a total rate: 6-17 years. 18-64 years. 65 years and older Total.
	The total is the sum of the age stratifications.

Two rates are reported:

- 1 Discharges for which the member received follow-up within 30 days after discharge
- 2 Discharges for which the member received follow-up within7 days after discharge



(FUH) Follow-Up After Hospitalization for Mental Illness (continued)

Description	Codes*
Outpatient Visit with a Mental Health Provider	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99492-99494, 99510, 99483 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
Visit Setting Unspecified Value Set with Partial Hospitalization POS with Mental Health Provider	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 52
Partial Hospitalization/Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Visit Setting Unspecified Value Set with Community Mental Health Center POS	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 53



(FUH) Follow-Up After Hospitalization for Mental Illness (continued)

Description	Codes*
Electroconvulsive Therapy with Ambulatory Surgical Center POS/ Community Mental Health Center POS/ Outpatient POS/Partial Hospitalization POS	CPT: 90870 Ambulatory POS: 24 Comm. POS: 53 Partial Hosp. POS: 52 Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
Telehealth Visit	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 02, 10
Observation	CPT: 99217-99220
Transitional Care Management	CPT: 99495, 99496
Telephone Visit	CPT: 98966-98968, 99441-99443
Psychiatric Collaborative Care Management	CPT: 99492-99494 HCPCS: G0512

^{*}Codes subject to change



(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder

Measure evaluates percentage of acute inpatient hospitalizations, residential treatment or withdrawal management visits for a diagnosis of substance use disorder among members 13 years of age and older that result in a follow-up visit or service for substance use disorder.

Two rates are reported:

- 1 The percentage of visits or discharges for which the member received follow-up for substance use disorder within the 30 days after the visit or discharge.
- 2 The percentage of visits or discharges for which the member received follow-up for substance use disorder within the 7 days after the visit or discharge.

Note: Follow-up does not include withdrawal management.

Description

Codes*

An Acute or Nonacute Inpatient Admission or Residential Behavioral Health Stay with a Principal Diagnosis of SUD on the Discharge Claim F10.xx-F19.xx



(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder (continued)

Description	Codes*
Outpatient Visit with a Principal Diagnosis of SUD	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 994883, 99492 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 ICD-10: F10.xx-F19.xx POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72
Intensive Outpatient Encounter or Partial Hospitalization with a Principal Diagnosis of SUD	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 HCPCS: GO410, GO411, HO035, H2001, H2012, S0201, S9480, S9484, S9485 ICD-10: F10.xx-F19.xx POS: 52

(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder (continued)

Description	Codes*
Non-residential Substance Abuse Treatment Facility with a Principal Diagnosis of SUD	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx POS: 57, 58
Community Mental Health Center Visit with a Principal Diagnosis of SUD	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx POS: 53
Telehealth Visit with a Principal Diagnosis of SUD	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx POS: 02, 10
Substance use Disorder Services with a Principal Diagnosis of SUD	CPT: 99408, 99409 HCPCS: : T1012, G0396, G0397, H0001, H0005, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, H0006, H0028
Opioid Treatment Service that Bills Monthly or Weekly with a Principal Diagnosis of SUD	HCPCS: G2086, G2087, G2071, G8074-G2077, G2080 ICD-10: F10.xx-F19.xx
Observation Visit with a Principal Diagnosis of SUD	CPT: 99217, 99218, 99219, 99220 ICD-10: F10.xx-F19.xx



(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder (continued)

Description	Codes*
Residential Behavioral Health Treatment with a Principal Diagnosis of SUD	HCPCS: H0017, H0018, H0019, T2048 ICD-10: F10.xx-F19.xx
Telephone Visit with a Principal Diagnosis of SUD	CPT: 98966-98968, 99441-99443 ICD-10: F10.xx-F19.xx
E-Visit or Virtual Check in with a Principal Diagnosis of SUD	CPT: 98969-98972, 99421-99444, 99457, 99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252
Pharmacotherapy Dispensing Event or Medication Treatment Event	Medications: Disulfiram (oral), Naltrexone (oral and injectable), Acamprosate (oral; delayed-release tablet), Buprenorphine (implant, injection, or sublingual tablet), Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) HCPCS: G2069, G2070, G2072, G2073, H0020, H0033, J0570-J0575, J2315, Q9991, Q9992, S0109

^{*}Codes subject to change



(FUM) Follow-Up After Emergency **Department Visit for Mental Illness**

Applicable Foster Care Measure:

Measure evaluates the percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness.

Two rates are reported:

- The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days).
- The percentage of ED visits for which the member received follow-up within 7 days of the ED visit (8 total days).

Description

Outpatient Visit with a Principal Diagnosis of a Mental Health Disorder

Codes*

CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211- 99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 994883, 99492 **HCPCS:** G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 ICD-10: F10.xx-F99

POS: 03, 05, 07, 09, 11-20, 22, 33,

49, 50, 71-72



(FUM) Follow-Up After Emergency Department Visit for Mental Illness (continued)

Description	Codes*
Intensive Outpatient Encounter or Partial Hospitalization with a Principal Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 ICD-10: F10.xx-F99 POS: 52
Community Mental Health Center Visit with a Principal Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F99 POS: 53
Electroconvulsive Therapy with a Principal Diagnosis of a Mental Health Disorder	CPT: 90780 POS: 03, 05, 07, 09, 11- 20, 22, 24, 33, 49, 50, 52, 53, 71, 72
Telehealth Visit with a Principal Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F99 POS: 02, 10
Observation Visit with a Principal Diagnosis of a Mental Health Disorder	CPT: 99217, 99218, 99219, 99220 ICD-10: F10.xx-F99
Telephone Visit with a Principal Diagnosis of a Mental Health Disorder	CPT: 98966-98968, 99441-99443 ICD-10: F10.xx-F99



(FUM) Follow-Up After Emergency Department Visit for Mental Illness (continued)

Description	Codes*
E-Visit or Virtual Check in with a Principal Diagnosis of a Mental Health Disorder	CPT: 98969-98972, 99421-99444, 99457, 99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252
An Outpatient with a Principal Diagnosis of Intentional Self-Harm with any Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 994883, 99492 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99 POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72
Intensive Outpatient Encounter or Partial Hospitalization with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99 POS: 52



(FUM) Follow-Up After Emergency Department Visit for Mental Illness (continued)

Description	Codes*
Community Mental Health Center Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99 POS: 53
Electroconvulsive Therapy with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder	CPT: 90780 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99 POS: 03, 05, 07, 09, 11- 20, 22, 24, 33, 49, 50, 52, 53, 71, 72
Telehealth Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99 POS: 02, 10
Observation Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder	CPT: 99217, 99218, 99219, 99220 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99
Telephone Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder	CPT: 98966-98968, 99441-99443 ICD-10: T40.xxxx-T43.xxxx, T51. xxxx with F10.xx-F99



(FUM) Follow-Up After Emergency Department Visit for Mental Illness (continued)

Description	Codes*
E-Visit or Virtual Check In with a	CPT: 98969-98972, 99421-99444,
Principal Diagnosis of Intentional	99457, 99458
Self-harm with any Diagnosis of a	HCPCS: G0071, G2010, G2012,
Mental Health Disorder	G2061-G2063, G2250-G2252
	ICD-10: T40.xxxx-T43.xxxx, T51.
	xxxx with F10.xx-F99

^{*}Codes subject to change



(IET) Initiation and Engagement of Substance Use Disorder Treatment

Measure evaluates percentage of adolescent and adult members with a new episode of substance use disorder (SUD) episodes that result in treatment initiation and engagement.

Tips:

- Explain the importance of a follow-up to your patients.
- · Schedule an initial follow-up appointment within 14 days.
- Reschedule patients as soon as possible who do not keep initial appointments.
- $\boldsymbol{\cdot}$ Use telehealth where appropriate.
- · Submit applicable codes.
- ✓ Initiation of SUD Treatment: percentage of new SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment within 14 days
- ✓ Engagement of SUD Treatment: percentage of new SUD episodes that have evidence of treatment engagement within 34 days of initiation



(IET) Initiation and Engagement of Substance Use Disorder Treatment (continued)

Alcohol Use Disorder Treatment Medications		
Description Prescription		
Aldehyde Dehydrogenase Inhibitor	Disulfiram (oral)	
Antagonist	Naltrexone (oral and injectable)	
Other	Acamprosafe (oral; delayed-release tablet)	

Opioid Use Disorder Treatment Medications		
Description	Prescription	Medication Lists
Antagonist	Naltrexone (oral)	Naltrexone Oral Medication List
Antagonist	Naltrexone (injectable)	Naltrexone Injection Medication List
Partial Agonist	Buprenorphine (sublingual tablet)	Buprenorphine Oral Medication List
Partial Agonist	Buprenorphine (injection)	Buprenorphine Injection Medication List
Partial Agonist	Buprenorphine (implant)	Buprenorphine Implant Medication List
Partial Agonist	Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)	Buprenorphine Naloxone Medication List

Tips:

- $\boldsymbol{\cdot}$ Explain the importance of follow-up to your patients.
- $\cdot\,$ Schedule an initial follow-up appointment within 14 days.
- Reschedule patients as soon as possible who do not keep initial appointments.
- $\boldsymbol{\cdot}$ Use telehealth where appropriate.
- \cdot Submit applicable codes.



(IET) Initiation and Engagement of Substance Use Disorder Treatments (continued)

Description	Codes*
Initiation and Engagement/ Treatment	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99221-99223, 99231, 99232, 99233, 99238, 99239, 99241-99245, 99341-99345, 99347-99350, 99251-99255, 99381-99387, 99391-99397, 99401-99404, 99408, 99409, 99411, 99412, 99483, 99492-99494, 99510 HCPCS: G0155, G0176, G0177, G0396, G0397, G0409, G0443, G0463, G2086, G2087, G0512, G2067-G2078, G2080, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0022, H0031, H0034-H0037, H0039, H0040, H0047, H0050, H2000, H2010, H2011, H2013-H2020, H2035, H2036, S0201, S9480, S9484, S9485, T1006, T1012, T1015 POS: 02, 03, 05, 07, 09, 11-20, 22, 33, 49-50, 52-53, 57, 58, 71-72
Telephone Visits	CPT: 98966-98968, 99441-99443
E-visit/Virtual Check-In	CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063

^{*}Codes subject to change



(POD) Pharmacotherapy for Opioid Use Disorder

Evaluates the percentage of opioid use disorder (OUD) pharmacotherapy events that lasted at least 180 days among members 16 years of age and older with a diagnosis of OUD and a new OUD pharmacotherapy event.

Description	Codes*
Opioid Use Disorder (OUD)	F11.10, F11.120-122, F11.129, F11.13-14, F11.150-151, F11.159, F11.181-182, F11.188, F11.19-20, F11.220-222, F11.229, F11.23-24, F11.250-251, F11.259, F11.281-282, F11.288, F11.29
Description	Prescription
Antagonist	Naltrexone (oral or injectable)
Partial Agonist	Buprenorphine (sublingual tablet, injection, or implant)
Partial Agonist	Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)
Agonist	Methadone (oral, medical claim codes H0020, S10109, G2067, G2078)

^{*}Codes subject to change

Tips:

- Provide empathetic listening and nonjudgmental discussions to engage patient.
- · Create a relapse prevention plan.
- Inform patient of risks and benefits of pharmacotherapy, treatment without medication, and no treatment.
- Offer mutual help like peer recovery support, harm reduction, 12-step fellowships.
- Reach out proactively within 24 hours if the patient does not keep scheduled appointment.
- Encourage coordination of care between physical and behavioral health providers.
- · Submit applicable codes.





(SAA) Adherence to Antipsychotic Medications for Individuals With Schizophrenia

Evaluates percentage of members 18 years of age and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.

Oral Antipsychotics		
· Aripiprazole	·Lumateperone	· Chlorpromazine
· Asenapine	· Lurasidone	· Fluphenazine
· Brexpiprazole	 Molindone 	· Perphenazine
· Cariprazine	· Olanzapine	· Prochlorperazine
· Clozapine	· Paliperidone	· Thioridazine
· Haloperidol	· Quetiapine	· Trifluoperazine
· Iloperidone	· Risperidone	· Amitriptyline-perphenazine
· Loxapine	· Ziprasidone	· Thiothixene

Description	Long Acting Injections Prescription
Long-acting Injections 14 Days Supply	Risperidone (excluding Perseris®)
Long-acting Injections 28 Days Supply	

Long-acting Injections Risperidone (Perseris®)
30 days Supply





(SSD) Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications

Measure evaluates percentage of members 18-64 years of age with schizophrenia, schizoaffective disorder, or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test.

Tips:

- Provide members with HbA1c or glucose test orders to be completed yearly.
- Educate the member about the risks associated with taking antipsychotic medications and the importance of regular follow up care.
- · Submit applicable codes.

Description	Codes*
HbA1c Lab Tests	CPT: 83036, 83037 CPT-CAT-II: 3044F, 3046F, 3051F, 3052F
Glucose Lab Tests	CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951

^{*}Codes subject to change



(UOP) Use of Opioids from Multiple Providers

Assesses the percentage of members 18 years and older, receiving prescription opioids for ≥15 days during the measurement year, who received opioids from multiple providers.

Three rates reported:

- **Multiple Prescribers-** The percentage of members receiving prescriptions for opioids from four or more different prescribers during the measurement year.
- 2 Multiple Pharmacies The percentage of members receiving prescriptions for opioids from four or more different pharmacies during the measurement year.
- 3 Multiple Prescribers and Multiple Pharmacies— The percentage of members receiving prescriptions for opioids from four or more different prescribers and four or more different pharmacies during the measurement year (i.e., the percentage of members who are numerator compliant for both the Multiple Prescribers and Multiple Pharmacies rates).

Note: A lower rate indicates better performance for all three rates.

Opioid Medications		
· Benzhydrocodone	· Meperidine	
· Buprenorphine	· Methadone	
(transdermal patch and buccal film)	· Morphine	
· Codeine	· Opium	
· Dihydrocodeine	· Oxycodone	
· Fentanyl	· Oxymorphone	
· Hydrocodone	· Pentazocine	
· Hydromorphone	· Tapentadol	
· Levorphanol	· Tramadol	

(Opioid medications exclude injectables and opioid-containing cough and cold products)

*For the follow-up treatments, include an ICD-10 diagnosis for Alcohol or Other Drug Dependence from the Mental, Behavioral and Neurodevelopmental Disorder Section of ICD-10 along with a procedure code for the preventive service, evaluation, and management consultation or counseling service.









